



King's Research Portal

DOI:

[10.1111/cei.13351](https://doi.org/10.1111/cei.13351)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Mondelli, V., & Vernon, A. C. (2019). From early adversities to immune activation in psychiatric disorders: the role of the sympathetic nervous system. *Clinical and Experimental Immunology*, 197(3), 319-328.
<https://doi.org/10.1111/cei.13351>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

From early adversities to immune activation in psychiatric disorders: the role of the sympathetic nervous system

Valeria Mondelli^{1, 2} and Anthony C. Vernon^{3,4}

¹ King's College London, Institute of Psychiatry Psychology and Neuroscience,
Department of Psychological Medicine

² NIHR Biomedical Research Centre South London and Maudsley NHS Trust

³ King's College London, Institute of Psychiatry Psychology and Neuroscience,
Department of Basic and Clinical Neuroscience

⁴ MRC Centre for Neurodevelopmental Disorders, King's College London

Correspondence should be addressed to either:

Dr Valeria Mondelli

Maurice Wohl Clinical Neuroscience Institute,
Cutcombe Road, SE5 9RT, London, United Kingdom

Email: valeria.mondelli@kcl.ac.uk

Phone: +44(0)2078480353

Dr Anthony Vernon

Maurice Wohl Clinical Neuroscience Institute,
Cutcombe Road, SE5 9RT, London, United Kingdom

Email: anthony.vernon@kcl.ac.uk

Phone: +44(0)2078485311

Summary

Increased peripheral levels of cytokines and central microglial activation have been reported in patients with psychiatric disorders. The degree of both innate and adaptive immune activation is also associated with worse clinical outcomes and poor treatment response in these patients. Understanding the possible causes and mechanisms leading to this immune activation is therefore an important and necessary step for the development of novel and more effective treatment strategies for these patients. In this manuscript, we review the evidence of literature pointing to childhood trauma as one of the main causes behind the increased immune activation in patients with psychiatric disorders. We then discuss the potential mechanisms linking the experience of early-life adversity (ELA) to innate immune activation. Specifically, we focus on the innervation of the bone marrow from sympathetic nervous system (SNS) as new and emerging mechanism that has the potential to bridge the observed increases in both central and peripheral inflammatory markers in patients exposed to ELA. Experimental studies in laboratory rodents suggest that SNS activation following early-life stress exposure causes a shift in the profile of innate immune cells, with an increase in pro-inflammatory monocytes. In turn, these cells traffic to the brain and influence neural circuitry, which manifests as increased anxiety and other relevant-behavioral phenotypes. To date however, very few studies have been conducted to explore this candidate mechanism in humans. Future research is also needed to clarify whether these pathways could be partially reversible to improve prevention and treatment strategies in the future.

Key words: childhood trauma, early adversity, inflammation, cytokine, microglia, sympathetic nervous system, depression, psychosis.

1. Introduction

In the past two decades, converging lines of evidence from several studies suggest a role for activation and/or dysfunction of the immune system in the pathogenesis of major psychiatric disorders such as depression and psychotic disorders. Specifically, there is evidence for increased peripheral immune activation, typically characterized by elevations in the levels of circulating pro-inflammatory cytokines such as interleukin (IL)-6 or IL-1 β . In addition, other inflammatory markers, such as C reactive protein (CRP) are also commonly increased in these patients. Finally, in both depression and psychotic disorder there is evidence from *post-mortem* brain tissue and *in vivo* neuroimaging studies to suggest central microglia activation, the brain's resident innate immune cell, although these data are not unequivocal (1, 2). The increased innate immune activation in these patients may be of particular clinical relevance, based on the findings of our group and others that the degree of immune activation, at least in terms of peripheral cytokines and CRP, is associated with more severe clinical symptoms, suicidal ideation and poor response to treatment (2-4).

In order to develop novel treatment and/or preventative measures for these patients, we need to better understand what are the causes and potential mechanisms underlying the immune activation in patients with psychiatric disorders. In this review we seek to address this question by first reviewing the evidence that links the experience of traumatic and stressful events during childhood to innate immune activation and development of psychopathology in later life (1). Childhood trauma is, unfortunately, highly prevalent in our society and one of the most widely recognized risk factors for development of psychiatric disorders later in life, including major depression, psychosis, and post-traumatic stress disorder (PTSD) (5-7). Research carried out in the past two decades has clearly identified a role for the activation of the immune system as a potential biological mechanism connecting childhood trauma and the subsequent development of psychopathology (8). In the second part of the

manuscript, we focus on the emerging evidence for a role of the sympathetic nervous system (SNS) as one of the main mechanisms behind the link between early life adversities (ELA) and peripheral and central inflammation. With this critical overview, the paper shows and identifies necessary next steps for research in this field, to enable the future development of clinical preventative and treatment strategies in victims of childhood trauma.

2. Evidence for the association between early life adversities and activation of the innate immune system

2.1 Increased circulating cytokines and acute phase proteins

The presence of increased immune activation in patients with psychiatric disorders across different diagnostic categories has generated increasing interest in unraveling causes behind this immune activation. In the context of peripheral immune activation, several clinical studies so far have shown that inflammatory markers, such as IL-6 and C reactive protein (CRP), are particularly high in patients who had experienced childhood traumatic events including physical or sexual abuse, emotional abuse or neglect that occur in childhood; this has been demonstrated both in patients suffering with psychosis, major depression or bipolar disorder (9-13). These primary research findings are supported by evidence from systematic reviews that provide evidence for prolonged peripheral inflammation following exposure to traumatic events in childhood (8, 11, 14). Increased levels of other peripheral circulating inflammatory markers, such as the acute phase protein CRP, are also widely reported in young children exposed to adverse psychosocial events (15, 16). Moreover, this initial early immune activation appears to persist and thus become chronic throughout adult life (14). In a previous meta-analysis, we reported evidence to suggest that adult subjects with experience of childhood trauma have higher levels not only of CRP, but also of the major pro-inflammatory cytokines interleukin (IL)-6 and tumor necrosis factor (TNF)- α , when compared with adults without such experience (8).

One important point in this regard is that the magnitude of the increases in these inflammatory markers is not comparable to that found following acute infections. Rather, it more closely resembles a low-threshold, chronic inflammation, which is broadly comparable to detectable levels of these markers in patients with cardiovascular disease or rheumatoid arthritis. Although modest in terms of intensity, this persistent immune activation appears to be relevant for development of psychiatric disorders later on in life. Specifically, data derived from longitudinal cohorts of children provide evidence for statistically significant associations between experiencing childhood traumatic events and increases in peripheral CRP levels, which are particularly pronounced in those individuals who later developed depression in adult life (17, 18).

2.2 Circulating immune cell populations – evidence for changes in both innate and adaptive immunity following ELA exposure

In contrast to the wealth of data on circulating cytokines following early life trauma exposure, proportionally far fewer studies have directly investigated the effects of exposure to childhood adversities on circulating immune cell populations using for example, fluorescence activated cell sorting (FACS). In adolescents who experienced early life stress compared to those without such experience, do Prado and colleagues reported increases in circulating T-cells expressing the activation markers CD2⁺/CD4⁺/CD25⁺ and CD3⁺/CD69⁺, but also an increase in senescent T cells (CD8⁺/CD28⁻ and CD4⁺/CD28⁻ cells) (19). These changes were further accompanied by decreased percentages of Natural Killer (NK; CD3⁻/CD56⁺) and NK T-cells (CD3⁺/CD56⁺) (19). Comparing individuals separated from their mothers in early life to those reared by their parents, Elwenspoek and colleagues also reported increased T-cell senescence (CD57⁺ cells) (20). In another FACS study from this group, comparing

individuals with a history of ELA with those without, a reduction in the numbers of CD69⁺/CD8⁺ T-cells and increased numbers of HLA-DR⁺/CD4⁺, HLA-DR⁺/CD8⁺ and CD25⁺/CD8⁺ T-cells was reported (21). Collectively, these results suggest that ELA exposure is associated with a sustained activation of adaptive immunity and accelerated immune ageing, as indexed by the increase in T-cell senescence. These effects appeared to be independent of either stress or health-risk behaviours, suggestive of a primary effect of early-life programming on specific immune cell populations (20-22). Exploring how social adversity may affect peripheral immune cells, Counotte et al. (2018) observed that individuals with high clinical risk for psychosis who were exposed to ELA were characterized by an increased number of CD4⁺ T-helper (Th) 17 cells when compared with those at lower risk (23). There were also significant predictive relationships between increased numbers of T regulator (T-reg) cells and decreased NK cell numbers, with the increased degree of stress experienced during exposure to virtual social stressors (23). If and how these different associations fit together still needs to be elucidated. Other studies characterizing the effects of social adversities on peripheral leukocyte populations have mainly identified increases in circulating monocytes and dendritic cells following exposure to traumatic experiences in early life (24, 25). Furthermore, Schwaiger and colleagues identified an altered stress-induced gene expression in CD14⁺ monocytes of adults with history of childhood trauma; the differences were particularly linked to the expression of genes involved in steroid hormone activity and signal transduction which were in turn associated with increased activity of pro-inflammatory signaling upstream (26).

These clinical data are supported by a large body of evidence from pre-clinical studies in adult mice exposed to repeated social defeat (RSD), which results in a significant accumulation of neutrophils and CD11b⁺ LyC6^{Hi} monocytes in both the circulation and spleen (27, 28). Splenic dendritic cells (DCs) from mice exposed to RSD also have increased surface expression of MHC-I, CD80 and CD44 indicative of an activated

state (29). It should also be noted however, that CD44 is commonly used as a T cell activation marker (29). Furthermore, following *in vitro* stimulation, these cells display an up-regulated pro-inflammatory gene expression signature and produce comparatively higher levels of pro- and anti-inflammatory cytokines including IL-6, TNF- α and IL-10 (29). Extending these findings, exposure to RSD in mice and low socioeconomic status (SES) in humans, results in a relative expansion of a transcriptional profile associated with immature pro-inflammatory monocytes in peripheral blood mononuclear cells (PMBCs) (30). This is characterized (at least in the mouse) by increased activity of transcription factors that regulate both differentiation to the myeloid lineage but also pro-inflammatory effector functions of these cells, such as PU.1 (30). In addition, exposure to RSD in mice results in increased formation of Ly6C^{High} monocytes, which could be prevented using β -adrenoreceptor antagonists (30). Collectively, these data suggest a role for the sympathetic nervous system driving, at least in part, up-regulation of myelopoiesis, characterized by a pro-inflammatory state, which we will return to later in this review (30). Data on the effects of social stress on adaptive immunity in pre-clinical models is less developed, although there are data suggesting a reduction in the numbers of T-cells in the circulation and spleen (27, 28), which is at odds with some of the aforementioned clinical findings. On the other hand, RSD is reported to result in enhanced CD4⁺ and CD8⁺ memory T cell responses when this stress is experienced before a viral infection (31-33). These data suggest a view that exposure to RSD in adult mice may significantly enhance virus-specific immune memory through the augmentation of specific memory T cell populations (31-33). Interestingly, using a mediation analysis, Elwenspoek and colleagues found that cytomegalovirus (CMV) infection largely explained the increase in T-cell senescence markers (CD57⁺) following ELA exposure, suggesting that viral infections may affect T-cell populations in humans following ELA exposure (20). When comparing these data it should however be remembered that the great majority of pre-

clinical studies that have characterized the peripheral immune response to stress exposure, have primarily focused on the effects of repeated social defeat stress (RSD) exposure in *adult* animals. Detailed peripheral immunophenotyping data regarding the specific effects of early-life stress in pre-clinical models is however, currently lacking. Another important point is that the effects of any experimental stress paradigm in animals may have both common and distinct effects on the immune system depending on the nature and duration of the stressor (34-36). In this context however, pre-clinical studies could be immensely valuable, since they allow the experimenter to probe the effects of different types of early-life stress with a precise control of genetic and other environmental variables, something that is very challenging in humans. Pre-clinical studies therefore have the clear potential to help dissect the specific peripheral immune cell responses associated with exposure to different stressors at different vulnerable periods from the prenatal period, through early postnatal life and into adolescence.

Overall, the evidence from the few studies available suggests that ELA exposure is associated with (1) direct programming of innate and adaptive immune cell function; (2) increased numbers of circulating innate and adaptive immune cells with a pro-inflammatory phenotype, a finding consistent with the elevation in circulating pro-inflammatory cytokines and acute phase proteins in those exposed to ELA and (3) accelerated immune ageing as indicated by T-cell senescence. What remains unclear from these data however is whether these peripheral changes are related to central immune activation. Here again, studies pre-clinical models of stress exposure are vital, since they permit invasive assessments in both the brain and periphery to aid dissection of the mechanisms governing immune-to-brain communication and vice-versa (37, 38). However, in addressing this question, we must first consider the clinical evidence base for central immune activation as a function of ELA exposure.

2.3 Central immune activation following ELA exposure

Assessment of central immune activation in clinical samples is typically achieved through *post-mortem* neuropathological assessment at either the cellular or molecular level, or *in vivo*, using positron emission tomography (PET) combined with selective radioligands for the translocator protein (TSPO) comparing patient groups with healthy age/sex matched controls (37). Searching the extant literature, we found only one TSPO PET study which had tested for an association between psychosocial risk factors for psychosis, including childhood trauma and central TSPO radioligand binding. Dahoun and colleagues measured TSPO binding in a small sample of healthy individuals who were classified into two groups on the basis of exposure to psychosocial risk (high-risk group; n=12) or no significant exposure to psychosocial risk (low-risk group; n=12) (39). There were however no significant group differences however in TSPO volume of distribution (V_T) in either total gray matter or the frontal, parietal, temporal and occipital lobes (39). At first glance, these data are not consistent with the hypothesis that exposure to stress and/or ELA is associated with increased central microglial activation. It should be noted however, that these were healthy individuals without a psychiatric diagnosis, raising this possibility that the “high-risk” group in this case are a resilient group in whom putative central microglial activation may not be seen (39). Moreover, although some of the stressors assessed in these individuals occurred recently there was a temporal separation between stress exposure and TSPO PET imaging. Hence, it remains possible that acute stress exposure could be associated with increased TSPO, which may normalize over time (39). Further studies are clearly required to address these issues and explore in more detail the effect of ELA on central TSPO binding in both pre-clinical models and clinical populations. In contrast, several TSPO PET studies have been conducted to search for any evidence of central immune activation in the psychiatric disorders for which ELA is a known risk factor, such as schizophrenia and depression. The data published so far suggest that there is evidence for putative central innate immune activation

associated with psychiatric disorders for which ELA exposure is a known risk factor (37), but evidence for specific effects of ELA on central immune cells in humans is currently lacking. Hence, to address this question, we must consider the evidence from pre-clinical rodent models of ELA exposure.

Exposing rodents or non-human primates to early life stress increases anxiety-like behaviors and impairs cognitive function in adulthood, suggesting that animal models may provide important insights into the effects of ELA in humans (40-42). As mentioned already, it should be remembered when comparing data from pre-clinical models to clinical studies that the effects of any experimental stress paradigm in animals may have both common and distinct effects on the immune system depending on the nature and duration of the stressor. This is no less true for central immune activation following stress exposure in rodents (36, 37). Interestingly, genetically controlled depletion of brain microglia in healthy rodents in early life is associated with abnormal social and motivated behaviour in a sex-specific manner, strongly implicating that microglia play a role in the early life programming of behaviour (43, 44). With regard to the evidence that early life stress exposure influences central microglia, several pre-clinical studies in different early-life stress models provide converging lines of evidence for increased innate immune activation in the central nervous system (CNS). Prenatal stress exposure, induced by exposing pregnant rat dams to restraint or other physical stressors, is reported to increase the number of Iba1⁺/CD68⁺ microglia, coupled with a shift their morphology to an activated, amoeboid state and decreased process motility, as well as increase pro-inflammatory cytokine expression in the brains of the affected offspring early in post-natal life primarily (but not exclusively) in the hippocampus, amygdala and pre-frontal cortex (45-54). None of these studies however, have investigated the effects of ELA exposure on TSPO binding, limiting cross-species comparisons to humans. At the molecular level,

Delpech and colleagues have examined the effect of brief daily separation (BDS) stress on the microglial transcriptome in the affected offspring, providing evidence that this early life stress actually perturbs the normal functional development of microglia in the developing hippocampus, with these cells displaying increased phagocytic activity and reduced expression of genes that normally increase across development (55). It is increasingly appreciated that microglia cells play critical roles in shaping the process of synaptic maturation as well as axonal guidance in the developing and maturing brain (56-64). Hence, any stress-induced perturbation of microglial physiology during these critical developmental windows would be predicted to result in the defective maturation of neural circuits (56, 57, 65). This could have long-lasting consequences on synaptic function and may be another mechanism whereby activation of central innate immunity after ELA exposure may directly increase the vulnerability to develop psychopathology in later life. These cells however may also play a role in central to peripheral immune communication as we discuss in the next section.

3. How does ELA lead to immune activation?

Although both clinical and pre-clinical studies provide converging lines evidence for both central and peripheral immune activation following ELA exposure (notably for both innate and adaptive immunity), a critical question remains as to how does ELA exposure specifically trigger both central and peripheral inflammation? Although the precise mechanisms by which peripheral immunity may influence central immunity and neuronal function (and vice versa) remain largely unknown, a number of interesting candidate mechanisms have been advanced. These include, 1) the development of glucocorticoid resistance and therefore an impairment in the anti-inflammatory action of glucocorticoids; 2) the activation of the sympathetic nervous system (SNS); 3) a change in the composition of gut microbiota, 4) exposure to physical injuries or infection as part of the early life trauma and 5) behavioral mechanisms, including

obesity, alcohol consumption and poor sleep which may themselves promote immune activation (66). For the purposes of this concise review, we will focus on the link between SNS activation in response to stress exposure and immune activation, for which there is a recent increasing body of evidence (67).

3.1 The role of the sympathetic nervous system in the link between early adversities and immune activation

The sympathetic nervous system (SNS) is one of the main biological systems involved in the stress response. Its activation also results in cascade of signals, which affect the regulation of the innate immune system, which occurs in cell-state and organ-specific manner. Of particular interest however is the bone marrow (BM), which is densely innervated by the SNS (68). The activation of the SNS, in response to stress, for example, leads to the production and release of norepinephrine (NE), which binds to β -adrenergic (β AR) and α -adrenergic (α AR) receptors expressed on peripheral immune cell populations in both rodents and humans (68, 69). Pre-clinical studies in animals show that SNS activation and NE production triggers the release of monocytes into the circulation, which can be modulated pharmacologically by systemic administration of either a β_2 AR agonist (increase) or β AR antagonist (decrease) (27, 30, 32). Increased numbers of circulating monocytes is also correlated with oscillations in SNS tone as a function of circadian rhythm (70) and can occur following other forms of SNS stimuli, such as exercise (71). These observations are also true for lymphocytes, with exposure to acute stress in humans resulting in mobilization of total monocytes and lymphocytes and CD8⁺ T lymphocytes (71). Nor-epinephrine (NE) released during SNS activation may however lead to different immunomodulatory effects, depending on the cell type, state and organ involved. For example, Case and Zimmerman (2016) propose a model in which NE release can either have a suppressive effect on naïve T-lymphocytes in lymphoid organs such as the spleen, but in other regions, such as the vasculature and renal system, NE may drive the activation

of pro-inflammatory Th17 T-lymphocytes (72).

Taken together, the aforementioned lines of evidence clearly show that SNS activation is well placed to mediate both innate and adaptive immunity and brain-immune communication, across species. There are however, no studies in healthy humans, either alone or in comparison to clinical populations, that provide direct evidence for a link between ELA exposure, SNS activation and elevated peripheral and central immunity, beyond the observations that stress-exposed individuals who have developed psychopathology have elevated numbers of circulating pro-inflammatory monocytes and lymphocytes – as already evidenced in this review. Here again, we must therefore turn to pre-clinical studies in rodent models to dissect this mechanism.

As already stated, the SNS innervates peripheral lymphoid tissues including the spleen and bone marrow. Pre-clinical data suggest that chronic stress exposure is associated with a decrease in expression of the chemokine CXCL12 in the hematopoietic stem cell niche, which in turn accelerates proliferation of hematopoietic stem cells resulting in enhanced production of neutrophils and monocytes (73). Increases in SNS outflow and NE release following RDS exposure also drives mobilization of lymphocytes from the murine spleen and bone marrow, which then enter the circulation (27, 28, 30, 32, 71, 74, 75). These data are consistent with the aforementioned findings in humans of increased number of leukocytes in the circulation of individuals with stress-related psychiatric disorders (76) and the aforementioned studies reported increases in these cells in the circulation of individuals exposed to ELA. The specific contribution of the rodent models however has been to demonstrate elegantly a potential connection between central and peripheral immunity, which highlights the involvement of the SNS. Specifically, RDS in *adult* mice, increases neural activation, (indexed by increased immunostaining for the immediate early gene c-Fos), in mouse brain regions that regulate fear and threat sensing, such as the amygdala, which in turn is associated

with both social avoidance and anxiety-like behaviour, that is sensitive to β AR antagonism, suggesting SNS involvement downstream of the activation of neural stress circuitry (75). In parallel, to this increased neural activity, RDS results in central microglial activation, with increased density of amoeboid microglia in the amygdala, PFC and hippocampus, with higher surface expression of CD68, CD14 and TLR-4, and increased expression of both IL-1 β and glucocorticoid responsive genes (e.g. FKBP5) in these microglia, indicative of activation (67, 74, 75, 77, 78). Moreover, CD11b⁺CD45^{High}Ly6C^{High} macrophages increased in the circulation, but also trafficked into the mouse brain, where they express higher amounts of IL-1 β and adopt a pro-inflammatory phenotype (see Figure 1) (67, 74, 75, 77, 78). These effects are dependent on β AR, since they may be blocked by propranolol (a β AR antagonist), suggestive of a role for SNS activation in the initial mobilization of the monocytes (75). Consistent with this view, the trafficking of monocytes from the spleen to the brain is necessary for the re-establishment of anxiety in stress-sensitized mice and this also occurs downstream of SNS activation (74, 79). Depletion of central microglia also prevents monocyte trafficking to the brain and blocks the development of anxiety behaviour developing following repeated RDS exposure (77). These data suggest a model in which stress exposure results in the elevated neural activity in stress circuits, leading to increased SNS outflow. The former triggers a microglial response, the latter triggers the mobilization of pro-inflammatory monocyte derived macrophages, which subsequently traffic to the brain in a CCR2 and IL-1 β -dependent manner involving communication between central microglia and the endothelial cells of the blood-brain barrier (38, 67, 80, 81). These elegant studies demonstrate a potential pathway by which SNS activation is a key part of a larger multi-faceted interaction between central neural activity in specific circuits, central microglial activation and mobilization of peripheral immunity, which then traffics to the brain to close the loop by influencing neural dynamics in the same circuits, promoting the emergence of anxiety-like

phenotypes.

It is worth emphasizing again here that this model has been developed based on experiments conducted in *adult* animals and in the context of the RDS paradigm. Hence, two important pre-clinical extensions of this work are necessary. First, to confirm whether this mechanism is necessary and sufficient for the development of abnormal behaviour (such as anxiety) following exposure to other types of stressor, including social isolation (SI), repeated social defeat stress (RSDS) or chronic unpredictable mild stress (CUMS). In this context, exposure to chronic variable stress is reported to increase Lin⁻Sca-1⁺c-Kit⁺ hematopoietic stem cells in the BM (73). Moreover, in mice susceptible to RSDS, elevated peripheral IL-6 and increased circulating pro-inflammatory Ly6C^{Hi} monocytes are reported, which also traffic to the perivascular space to increase central IL-6 leading to depressive-like behaviour (80). Lastly, as already described herein, there is also evidence for central microglial activation in several animal models of early-life stress, although forward translation of this evidence to humans using TSPO binding for example is currently lacking (39). These data may suggest that there are indeed common mechanisms across different stress exposures, but further work is needed in this respect to clarify this mechanistically. Second, studies in the pre-, perinatal and adolescent period are necessary to directly test if the model proposed from adult animals contributes to the association between ELA exposure and psychopathology in later life. It is worth recalling here that microglia depletion by itself in this period (without stress exposure) was sufficient to induce long-term emotional behavioral changes in mice (43). Moreover, molecular studies suggest that microglia development may be abnormal following ELA exposure, perhaps indicative of later loss-of-function, or reduced flexibility of these cells (55, 82). Hence, the relationships between neural circuit activity central and peripheral immunity may differ in the developing as compared to the adult brain, which requires investigation (82, 83).

From the clinical perspective, direct evidence for involvement of the SNS *per se* following ELA exposure, still needs to be provided. Indeed, the evidence supporting the link between childhood trauma, for example and hyper-activation of the SNS in humans is conflicting, with some studies reporting an increased reactivity of SNS following high levels of family adversities, while others finding no such association (84). For example, no study has so far investigated immune markers together with markers of SNS activation in individuals with experience of childhood trauma compared with relevant controls. This would be an extremely important step to understand if the immune activation is found mainly in those who present an enhanced reactivity of the SNS.

4. Conclusions

In conclusion, increasing evidence points towards a fundamental role of ELA exposure in the activation of both innate and adaptive arms of the immune system in patients with psychiatric disorders. The SNS appears to play a central role in this link by increasing the activation of T cells and shifting the differentiation of hematopoietic stem cells in the bone marrow to pro-inflammatory monocytes, with elevated expression of pro-inflammatory molecules. Pre-clinical studies in adult animals suggest trafficking of peripheral immune cells to the brain is an important mechanism for the development of anxiety following stress exposure. We highlight an important lack of studies addressing these issues, insofar as it is possible to do so in human samples, but also the lack of detailed work on these potential mechanisms which link central and peripheral immunity following early-life stress exposure in pre-clinical models. Future studies to address these questions have the potential to identify molecular targets and time windows of opportunity for both prevention and treatment strategies to ameliorate the devastating long-term effects of ELA on mental health.

Acknowledgments

Dr Mondelli is supported by MQ: Transforming Mental Health (Grant: MQBF1), the Medical Research Foundation (Grant: MRF-160-0005-ELP-MONDE), and the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health. Dr Vernon acknowledges the financial support of the Medical Research Council (New Investigator Research Grant MR/N025377/1 and Centre Grant MR/N026063/1). The authors are also supported by the King's Together Fund of King's College London.

Conflict of Interest

Dr Mondelli has received research funding from Johnson & Johnson as part of a research program on depression and inflammation. Dr Vernon has received research funding from F. Hoffman La Roche Ltd and UCB Biopharma SPRL as part of a research program on early life immune activation. The views expressed are those of the authors and not necessarily those of Johnson & Johnson, F Hoffman La Roche or UCB Biopharma SPRL. These funders had no influence on the decision to publish this work.

Legend to Figures

Figure 1: A schematic representation of how Early Life Adversities may contribute to peripheral and central immune activation through activation of the sympathetic nervous system (SNS). Orange arrows indicate neural pathways, blue indicate central microglial effects and green the influence of peripheral monocytes. **(1)** Exposure to early life adversity (ELA) such as childhood trauma results in **(2)** activation of the brain stress circuitry and **(3)** increased outflow of the sympathetic nervous system to the bone marrow with increased nor-epinephrine (NE) release. **(4)** This decreases the

expression of the chemokine CXCL12 in the bone marrow hematopoietic stem cell niche, which accelerates proliferation of hematopoietic stem cells resulting in enhanced production of pro-inflammatory CD14+/CD16- monocytes, which are released into circulation and **(5)** traffic to the blood-brain-barrier through the vasculature. **(6)** At the same time, central microglial cells are also activated in response to neural cues as a result of stress circuit activation. These cells facilitate the infiltration of the peripheral monocytes through secretion of specific cytokines and chemokines, which modulate endothelial cell permeability and allow ingress of the monocytes from the vasculature to the peri-vascular space and the brain parenchyma. **(7)** The infiltrating pro-inflammatory monocytes lead to further sustained activation of neural stress circuits, resulting in the emergence of abnormal behaviour. **(8)** In parallel, ELA exposure may also influence the development and function of central microglia. Converging lines of evidence suggest that microglia play key roles in shaping the development of the brain including the formation and pruning of dendritic spines and synapses. Hence, any disruption of this process following stress exposure in early life may result in the abnormal neural circuit development, which by itself may predispose an individual to psychopathology in later life. PFC: prefrontal cortex; AMYG: amygdala; HPC: hippocampus; NE: norepinephrine; SNS: sympathetic nervous system.

References

1. Baumeister D, Russell A, Pariante CM, Mondelli V. Inflammatory biomarker profiles of mental disorders and their relation to clinical, social and lifestyle factors. *Soc Psychiatry Psychiatr Epidemiol.* 2014;49(6):841-9.
2. Mondelli V, Vernon AC, Turkheimer F, Dazzan P, Pariante CM. Brain microglia in psychiatric disorders. *The lancet Psychiatry.* 2017.
3. Mondelli V, Ciufolini S, Belvederi Murri M, Bonaccorso S, Di Forti M, Giordano A, et al. Cortisol and Inflammatory Biomarkers Predict Poor Treatment Response in First Episode Psychosis. *Schizophr Bull.* 2015;41(5):1162-70.

4. Cattaneo A, Gennarelli M, Uher R, Breen G, Farmer A, Aitchison KJ, et al. Candidate genes expression profile associated with antidepressants response in the GENDEP study: differentiating between baseline 'predictors' and longitudinal 'targets'. *Neuropsychopharmacology*. 2013;38(3):377-85.
5. Schafer I, Fisher HL. Childhood trauma and psychosis - what is the evidence? *Dialogues Clin Neurosci*. 2011;13(3):360-5.
6. Infurna MR, Reichl C, Parzer P, Schimmenti A, Bifulco A, Kaess M. Associations between depression and specific childhood experiences of abuse and neglect: A meta-analysis. *J Affect Disord*. 2016;190:47-55.
7. Yehuda R, Hoge CW, McFarlane AC, Vermetten E, Lanius RA, Nievergelt CM, et al. Post-traumatic stress disorder. *Nat Rev Dis Primers*. 2015;1:15057.
8. Baumeister D, Akhtar R, Ciufolini S, Pariante CM, Mondelli V. Childhood trauma and adulthood inflammation: a meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor-alpha. *Mol Psychiatry*. 2016;21(5):642-9.
9. Aas M, Dieset I, Hope S, Hoseth E, Mørch R, Reponen E, et al. Childhood maltreatment severity is associated with elevated C-reactive protein and body mass index in adults with schizophrenia and bipolar diagnoses. *Brain Behav Immun*. 2017;65:342-9.
10. Di Nicola M, Cattaneo A, Hepgul N, Di Forti M, Aitchison KJ, Janiri L, et al. Serum and gene expression profile of cytokines in first-episode psychosis. *Brain Behav Immun*. 2013;31:90-5.
11. Hepgul N, Pariante CM, Dapasquale S, Diforti M, Taylor H, Marques TR, et al. Childhood maltreatment is associated with increased body mass index and increased C-reactive protein levels in first-episode psychosis patients. *Psychol Med*. 2012;1-9.
12. Grosse L, Ambree O, Jorgens S, Jawahar MC, Singhal G, Stacey D, et al. Cytokine levels in major depression are related to childhood trauma but not to recent stressors. *Psychoneuroendocrinology*. 2016;73:24-31.

13. Munjiza A, Kostic M, Pesic D, Gajic M, Markovic I, Tosevski DL. Higher concentration of interleukin 6 - A possible link between major depressive disorder and childhood abuse. *Psychiatry Res.* 2018;264:26-30.
14. Danese A, Pariante CM, Caspi A, Taylor A, Poulton R. Childhood maltreatment predicts adult inflammation in a life-course study. *Proc Natl Acad Sci U S A.* 2007;104(4):1319-24.
15. Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiol Behav.* 2012;106(1):29-39.
16. Schmeer KK, Yoon A. Socioeconomic status inequalities in low-grade inflammation during childhood. *Arch Dis Child.* 2016;101(11):1043-7.
17. Danese A, Moffitt TE, Pariante CM, Ambler A, Poulton R, Caspi A. Elevated inflammation levels in depressed adults with a history of childhood maltreatment. *Arch Gen Psychiatry.* 2008;65(4):409-15.
18. Khandaker GM, Zuber V, Rees JMB, Carvalho L, Mason AM, Foley CN, et al. Shared mechanisms between coronary heart disease and depression: findings from a large UK general population-based cohort. *Mol Psychiatry.* 2019.
19. do Prado CH, Grassi-Oliveira R, Daruy L, Wieck A, Bauer ME. Evidence for Immune Activation and Resistance to Glucocorticoids Following Childhood Maltreatment in Adolescents Without Psychopathology. *Neuropsychopharmacol.* 2017;42(11):2272-82.
20. Elwenspoek MMC, Sias K, Hengesch X, Schaan VK, Leenen FAD, Adams P, et al. T cell immunosenescence after early life adversity: association with cytomegalovirus infection. *Front Immunol.* 2017;8.
21. Elwenspoek MMC, Hengesch X, Leenen FAD, Schritz A, Sias K, Schaan VK, et al. Proinflammatory T Cell Status Associated with Early Life Adversity. *J Immunol.* 2017;199(12):4046-55.
22. Elwenspoek MMC, Kuehn A, Muller CP, Turner JD. The effects of early life adversity on the immune system. *Psychoneuroendocrino.* 2017;82:140-54.

23. Couston J, Drexhage HA, Wijkhuijs JM, Pot-Kolder R, Bergink V, Hoek HW, et al. Th17/T regulator cell balance and NK cell numbers in relation to psychosis liability and social stress reactivity. *Brain Behav Immun*. 2018;69:408-17.
24. Cole SW, Hawkley LC, Arevalo JM, Cacioppo JT. Transcript origin analysis identifies antigen-presenting cells as primary targets of socially regulated gene expression in leukocytes. *Proc Natl Acad Sci U S A*. 2011;108(7):3080-5.
25. O'Donovan A, Sun B, Cole S, Rempel H, Lenoci M, Pulliam L, et al. Transcriptional control of monocyte gene expression in post-traumatic stress disorder. *Dis Markers*. 2011;30(2-3):123-32.
26. Schwaiger M, Grinberg M, Moser D, Zang JC, Heinrichs M, Hengstler JG, et al. Altered Stress-Induced Regulation of Genes in Monocytes in Adults with a History of Childhood Adversity. *Neuropsychopharmacology*. 2016;41(10):2530-40.
27. Engler H, Dawils L, Hoves S, Kurth S, Stevenson JR, Schauenstein K, et al. Effects of social stress on blood leukocyte distribution: the role of alpha- and beta-adrenergic mechanisms. *J Neuroimmunol*. 2004;156(1-2):153-62.
28. Engler H, Bailey MT, Engler A, Sheridan JF. Effects of repeated social stress on leukocyte distribution in bone marrow, peripheral blood and spleen. *J Neuroimmunol*. 2004;148(1-2):106-15.
29. Powell ND, Bailey MT, Mays JW, Stiner-Jones LM, Hanke ML, Padgett DA, et al. Repeated social defeat activates dendritic cells and enhances Toll-like receptor dependent cytokine secretion. *Brain Behav Immun*. 2009;23(2):225-31.
30. Powell ND, Sloan EK, Bailey MT, Arevalo JMG, Miller GE, Chen E, et al. Social stress up-regulates inflammatory gene expression in the leukocyte transcriptome via beta-adrenergic induction of myelopoiesis. *P Natl Acad Sci USA*. 2013;110(41):16574-9.
31. Mays JW, Bailey MT, Hunzeker JT, Powell ND, Papenfuss T, Karlsson EA, et al. Influenza Virus-Specific Immunological Memory Is Enhanced by Repeated Social Defeat. *J Immunol*. 2010;184(4):2014-25.

32. Powell ND, Mays JW, Bailey MT, Hanke ML, Sheridan JF. Immunogenic dendritic cells primed by social defeat enhance adaptive immunity to influenza A virus. *Brain Behav Immun*. 2011;25(1):46-52.
33. Mays JW, Powell ND, Hunzeker JT, Hanke ML, Bailey MT, Sheridan JF. Stress and the anti-influenza immune response: Repeated social defeat augments clonal expansion of CD8(+)T cells during primary influenza A viral infection. *J Neuroimmunol*. 2012;243(1-2):34-42.
34. Alleva E, Santucci D. Psychosocial vs. "physical" stress situations in rodents and humans - Role of neurotrophins. *Physiol Behav*. 2001;73(3):313-20.
35. Sheridan JF, Padgett DA, Avitsur R, Marucha PT. Experimental models of stress and wound healing. *World J Surg*. 2004;28(3):327-30.
36. Calcia MA, Bonsall DR, Bloomfield PS, Selvaraj S, Barichello T, Howes OD. Stress and neuroinflammation: a systematic review of the effects of stress on microglia and the implications for mental illness. *Psychopharmacology*. 2016;233(9):1637-50.
37. Mondelli V, Vernon AC, Turkheimer F, Dazzan P, Pariante CM. Brain microglia in psychiatric disorders. *Lancet Psychiatry*. 2017;4(7):563-72.
38. Weber MD, Godbout JP, Sheridan JF. Repeated Social Defeat, Neuroinflammation, and Behavior: Monocytes Carry the Signal. *Neuropsychopharmacol*. 2017;42(1):46-61.
39. Dahoun T, Calcia MA, Veronese M, Bloomfield P, Reis Marques T, Turkheimer F, et al. The association of psychosocial risk factors for mental health with a brain marker altered by inflammation: A translocator protein (TSPO) PET imaging study. *Brain Behav Immun*. 2019.
40. Wei L, Hao J, Kaffman A. Early Life Stress Inhibits Expression of Ribosomal RNA in the Developing Hippocampus. *Plos One*. 2014;9(12).
41. Wei L, Hao J, Lacher RK, Abbott T, Chung L, Colangelo CM, et al. Early-Life Stress Perturbs Key Cellular Programs in the Developing Mouse Hippocampus. *Dev Neurosci-Basel*. 2015;37(6):476-88.

42. Wei L, Simen A, Mane S, Kaffman A. Early Life Stress Inhibits Expression of a Novel Innate Immune Pathway in the Developing Hippocampus. *Neuropsychopharmacol.* 2012;37(2):567-80.
43. Nelson LH, Lenz KM. Microglia depletion in early life programs persistent changes in social, mood-related, and locomotor behavior in male and female rats. *Behav Brain Res.* 2017;316:279-93.
44. Bilbo SD, Schwarz JM. The immune system and developmental programming of brain and behavior. *Front Neuroendocrinol.* 2012;33(3):267-86.
45. Gomez-Gonzalez B, Escobar A. Prenatal stress alters microglial development and distribution in postnatal rat brain. *Acta Neuropathol.* 2010;119(3):303-15.
46. Roque A, Ochoa-Zarzosa A, Torner L. Maternal separation activates microglial cells and induces an inflammatory response in the hippocampus of male rat pups, independently of hypothalamic and peripheral cytokine levels. *Brain Behav Immun.* 2016;55:39-48.
47. Delpech JC, Wei L, Hao J, Yu X, Madore C, Butovsky O, et al. Early life stress perturbs the maturation of microglia in the developing hippocampus. *Brain Behav Immun.* 2016;57:79-93.
48. Giovanoli S, Engler H, Engler A, Richetto J, Voget M, Willi R, et al. Stress in puberty unmasks latent neuropathological consequences of prenatal immune activation in mice. *Science.* 2013;339(6123):1095-9.
49. Giovanoli S, Engler H, Engler A, Richetto J, Feldon J, Riva MA, et al. Preventive effects of minocycline in a neurodevelopmental two-hit model with relevance to schizophrenia. *Transl Psychiatry.* 2016;6:e772.
50. Cohen S, Ke X, Liu Q, Fu Q, Majnik A, Lane R. Adverse early life environment increases hippocampal microglia abundance in conjunction with decreased neural stem cells in juvenile mice. *Int J Dev Neurosci.* 2016;55:56-65.
51. Wang HT, Huang FL, Hu ZL, Zhang WJ, Qiao XQ, Huang YQ, et al. Early-Life Social Isolation-Induced Depressive-Like Behavior in Rats Results in Microglial

Activation and Neuronal Histone Methylation that Are Mitigated by Minocycline. *Neurotox Res.* 2017;31(4):505-20.

52. Gong Y, Tong LJ, Yang RR, Hu WF, Xu XG, Wang WJ, et al. Dynamic changes in hippocampal microglia contribute to depressive like behavior induced by early social isolation. *Neuropharmacology.* 2018;135:223-33.

53. Reus GZ, Silva RH, de Moura AB, Presa JF, Abelaira HM, Abatti M, et al. Early Maternal Deprivation Induces Microglial Activation, Alters Glial Fibrillary Acidic Protein Immunoreactivity and Indoleamine 2,3-Dioxygenase during the Development of Offspring Rats. *Mol Neurobiol.* 2019;56(2):1096-108.

54. Chastain LG, Franklin T, Gangisetty O, Cabrera MA, Mukherjee S, Shrivastava P, et al. Early life alcohol exposure primes hypothalamic microglia to later-life hypersensitivity to immune stress: possible epigenetic mechanism. *Neuropsychopharmacol.* 2019.

55. Delpech JC, Wei L, Hao J, Yu XQ, Madore C, Butovsky O, et al. Early life stress perturbs the maturation of microglia in the developing hippocampus. *Brain Behav Immun.* 2016;57:79-93.

56. Paolicelli RC, Ferretti MT. Function and Dysfunction of Microglia during Brain Development: Consequences for Synapses and Neural Circuits. *Front Synaptic Neurosci.* 2017;9:9.

57. Zhan Y, Paolicelli RC, Sforazzini F, Weinhard L, Bolasco G, Pagani F, et al. Deficient neuron-microglia signaling results in impaired functional brain connectivity and social behavior. *Nat Neurosci.* 2014;17(3):400-6.

58. Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, et al. Synaptic pruning by microglia is necessary for normal brain development. *Science.* 2011;333(6048):1456-8.

59. Paolicelli RC, Gross CT. Microglia in development: linking brain wiring to brain environment. *Neuron Glia Biol.* 2011;7(1):77-83.

60. Schafer DP, Lehrman EK, Stevens B. The "quad-partite" synapse: microglia-synapse interactions in the developing and mature CNS. *Glia*. 2013;61(1):24-36.
61. Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, et al. Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron*. 2012;74(4):691-705.
62. Basilico B, Pagani F, Grimaldi A, Cortese B, Di Angelantonio S, Weinhard L, et al. Microglia shape presynaptic properties at developing glutamatergic synapses. *Glia*. 2019;67(1):53-67.
63. Weinhard L, di Bartolomei G, Bolasco G, Machado P, Schieber NL, Neniskyte U, et al. Microglia remodel synapses by presynaptic trogocytosis and spine head filopodia induction. *Nat Commun*. 2018;9(1):1228.
64. Weinhard L, Neniskyte U, Vadisiute A, di Bartolomei G, Aygun N, Riviere L, et al. Sexual dimorphism of microglia and synapses during mouse postnatal development. *Dev Neurobiol*. 2018;78(6):618-26.
65. Paolicelli RC, Gross CT. Microglia in development: linking brain wiring to brain environment. *Neuron Glia Biol*. 2011;7(1):77-83.
66. Danese A, S JL. Psychoneuroimmunology of Early-Life Stress: The Hidden Wounds of Childhood Trauma? *Neuropsychopharmacology*. 2017;42(1):99-114.
67. Wohleb ES, Delpech JC. Dynamic cross-talk between microglia and peripheral monocytes underlies stress-induced neuroinflammation and behavioral consequences. *Prog Neuro-Psychoph*. 2017;79:40-8.
68. Elenkov IJ, Wilder RL, Chrousos GP, Vizi ES. The sympathetic nerve--an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev*. 2000;52(4):595-638.
69. Gosain A, Muthu K, Gamelli RL, DiPietro LA. Norepinephrine suppresses wound macrophage phagocytic efficiency through alpha- and beta-adrenoreceptor dependent pathways. *Surgery*. 2007;142(2):170-9.

70. Curtis AM, Bellet MM, Sassone-Corsi P, O'Neill LA. Circadian clock proteins and immunity. *Immunity*. 2014;40(2):178-86.
71. Riddell NE, Burns VE, Wallace GR, Edwards KM, Drayson M, Redwine LS, et al. Progenitor cells are mobilized by acute psychological stress but not beta-adrenergic receptor agonist infusion. *Brain Behav Immun*. 2015;49:49-53.
72. Case AJ, Zimmerman MC. Sympathetic-mediated activation versus suppression of the immune system: consequences for hypertension. *J Physiol*. 2016;594(3):527-36.
73. Heidt T, Sager HB, Courties G, Dutta P, Iwamoto Y, Zaltsman A, et al. Chronic variable stress activates hematopoietic stem cells. *Nat Med*. 2014;20(7):754-8.
74. McKim DB, Patterson JM, Wohleb ES, Jarrett BL, Reader BF, Godbout JP, et al. Sympathetic Release of Splenic Monocytes Promotes Recurring Anxiety Following Repeated Social Defeat. *Biol Psychiat*. 2016;79(10):803-13.
75. Wohleb ES, Powell ND, Godbout JP, Sheridan JF. Stress-Induced Recruitment of Bone Marrow-Derived Monocytes to the Brain Promotes Anxiety-Like Behavior. *J Neurosci*. 2013;33(34):13820-+.
76. Beydoun MA, Beydoun HA, Dore GA, Canas JA, Fanelli-Kuczmarski MT, Evans MK, et al. White blood cell inflammatory markers are associated with depressive symptoms in a longitudinal study of urban adults. *Transl Psychiatry*. 2016;6(9):e895.
77. McKim DB, Weber MD, Niraula A, Sawicki CM, Liu X, Jarrett BL, et al. Microglial recruitment of IL-1 beta-producing monocytes to brain endothelium causes stress-induced anxiety. *Mol Psychiatr*. 2018;23(6):1421-31.
78. Wohleb ES, Mckim DB, Sheridan JF, Godbout JP. Monocyte trafficking to the brain with stress and inflammation: a novel axis of immune-to-brain communication that influences mood and behavior. *Front Neurosci-Switz*. 2015;8.
79. Wohleb ES, McKim DB, Shea DT, Powell ND, Tarr AJ, Sheridan JF, et al. Re-establishment of Anxiety in Stress-Sensitized Mice Is Caused by Monocyte Trafficking from the Spleen to the Brain. *Biol Psychiat*. 2014;75(12):970-81.

80. Menard C, Pfau ML, Hodes GE, Kana V, Wang VX, Bouchard S, et al. Social stress induces neurovascular pathology promoting depression. *Nat Neurosci.* 2017;20(12):1752-+.
81. Weber M, McKim D, Sawicki C, Niraula A, Godbout J, Sheridan J. Microglial Recruitment of Monocytes to the Brain Causes Stress-Induced Anxiety. *Neuropsychopharmacol.* 2016;41:S461-S2.
82. Johnson FK, Kaffman A. Early life stress perturbs the function of microglia in the developing rodent brain: New insights and future challenges. *Brain Behav Immun.* 2018;69:18-27.
83. Johnson FK, Delpech JC, Thompson GJ, Wei L, Hao J, Herman P, et al. Amygdala hyper-connectivity in a mouse model of unpredictable early life stress. *Transl Psychiatry.* 2018;8(1):49.
84. McLaughlin KA, Sheridan MA, Alves S, Mendes WB. Child maltreatment and autonomic nervous system reactivity: identifying dysregulated stress reactivity patterns by using the biopsychosocial model of challenge and threat. *Psychosom Med.* 2014;76(7):538-46.